

# Embeddability of infinite graphs

Robin Christian<sup>1</sup>   Bruce Richter<sup>2</sup>   Gelasio Salazar<sup>3</sup>

<sup>1</sup>Fachbereich Mathematik  
Universität Hamburg (Germany)

<sup>2</sup>Department of Combinatorics and Optimization  
University of Waterloo (Canada)

<sup>3</sup>Instituto de Física  
Universidad Autónoma de San Luis Potosí (Mexico)

Graph Theory @ Georgia Tech (RobinFest)  
May 7, 2012

# Embeddability in the plane: Kuratowski, Wagner

## Theorem (Kuratowski, 1930)

A **finite** graph  $G$  is embeddable in the plane if and only if it does not contain a subgraph homeomorphic to the complete graph  $K_5$  or the complete bipartite graph  $K_{3,3}$ .

# Embeddability in the plane: Kuratowski, Wagner

## Theorem (Kuratowski, 1930)

A **finite** graph  $G$  is embeddable in the plane if and only if it does not contain a subgraph homeomorphic to the complete graph  $K_5$  or the complete bipartite graph  $K_{3,3}$ .

## Theorem (Wagner, 1937)

A **finite** graph  $G$  is embeddable in the plane if and only if it contains neither  $K_{3,3}$  nor  $K_5$  as a minor.

# Embeddability in the plane: Kuratowski, Wagner

## Theorem (Kuratowski, 1930)

A **finite** graph  $G$  is embeddable in the plane if and only if it does not contain a subgraph homeomorphic to the complete graph  $K_5$  or the complete bipartite graph  $K_{3,3}$ .

## Theorem (Wagner, 1937)

A **finite** graph  $G$  is embeddable in the plane if and only if it contains neither  $K_{3,3}$  nor  $K_5$  as a minor.


Every compact surface has a “Wagner’s Theorem”:

## Theorem (Robertson and Seymour, 1990)

For every compact surface there is a finite list of graphs such that a graph  $G$  is embeddable in this surface if and only if it does not contain any of these as a minor.

# $g + 1$ disjoint Kuratowski graphs: natural obstacle for embedding in genus $g$

$K_{3,3}$  and  $K_5$  are the **Kuratowski graphs**.




Torus (genus 1): can host **one** Kuratowski graph (and no more).

The compact surface of genus  $g$  can host the disjoint union of  $g$  Kuratowski graphs (and no more)

# $g + 1$ disjoint Kuratowski graphs: natural obstacle for embedding in genus $g$

$K_{3,3}$  and  $K_5$  are the **Kuratowski graphs**.




Torus (genus 1): can host **one** Kuratowski graph (and no more).

The compact surface of genus  $g$  can host the disjoint union of  $g$  Kuratowski graphs (and no more)

$\implies$  having many disjoint (or “sufficiently” disjoint) Kuratowski graphs makes the genus grow

# $g + 1$ disjoint Kuratowski graphs: natural obstacle for embedding in genus $g$

$K_{3,3}$  and  $K_5$  are the **Kuratowski graphs**.



Torus (genus 1): can host **one** Kuratowski graph (and no more).

The compact surface of genus  $g$  can host the disjoint union of  $g$  Kuratowski graphs (and no more)

$\implies$  having many disjoint (or “sufficiently” disjoint) Kuratowski graphs makes the genus grow

Is this **the** reason why the genus of a graph grows?

# Why does the genus of a graph grow?



# Why does the genus of a graph grow?

## Robertson and Seymour (unpublished)

There is a function  $f(g)$  tending to infinity so that, if a graph  $G$  does not embed in any surface of Euler characteristic at least  $2 - 2g$ , then  $G$  has one of the following graphs as a minor:

- ①  $f(g)$  disjoint copies of either  $K_{3,3}$  or  $K_5$ ;
- ②  $f(g)$  copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for a common vertex;
- ③  $f(g)$  copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for two common vertices; or
- ④  $K_{3,f(g)}$

# Why does the genus of a graph grow?

## Robertson and Seymour (unpublished)

There is a function  $f(g)$  tending to infinity so that, if a graph  $G$  does not embed in any surface of Euler characteristic at least  $2 - 2g$ , then  $G$  has one of the following graphs as a minor:

- 1  $f(g)$  disjoint copies of either  $K_{3,3}$  or  $K_5$ ;
- 2  $f(g)$  copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for a common vertex;
- 3  $f(g)$  copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for two common vertices; or
- 4  $K_{3,f(g)}$

If we restrict ourselves to orientable surfaces, then we have to add the  $f(g)$ -projective grid to the list.

# Our problem

Which graphs do not embed into any surface of bounded genus?

# Our problem

Which graphs do not embed into any surface of bounded genus?

Plausible answer, in view of the Robertson-Seymour result

Those that contain as a minor either:

- Infinitely many “sufficiently disjoint”  $K_{3,3}$ ’s or  $K_5$ ’s.
- $K_{3,\infty}$  (yes, abuse of notation)

## Some previous fine-tuning

Which graphs do not embed into any surface of bounded genus?

## Some previous fine-tuning

Which graphs do not embed into any surface of bounded genus?

- Obviously, every finite graph embeds in some surface. So the question is interesting only for infinite graphs.

# Some previous fine-tuning

Which graphs do not embed into any surface of bounded genus?

- Obviously, every finite graph embeds in some surface. So the question is interesting only for infinite graphs.
- If an infinite graph embeds in a surface, then it has only countably many vertices of degree  $\geq 3$  (Wagner, 1967). Thus it consists of a countable graph, plus possibly continuumly many cycles, paths, rays, and double rays.

$\implies$  the focus is on *countably infinite* graphs

# Some previous fine-tuning

Which graphs do not embed into any surface of bounded genus?

- Obviously, every finite graph embeds in some surface. So the question is interesting only for infinite graphs.
- If an infinite graph embeds in a surface, then it has only countably many vertices of degree  $\geq 3$  (Wagner, 1967). Thus it consists of a countable graph, plus possibly continuumly many cycles, paths, rays, and double rays.

$\implies$  the focus is on *countably infinite* graphs

Thus, it all boils down to:

## Question

Which countable graphs embed in **some** surface of bounded genus?



# Our result: embeddability in bounded genus

Theorem (Christian, Richter, and S., 2011<sup>+</sup>)

A countable graph  $G$  embeds in some (orientable) surface of bounded genus if and only if  $G$  does not contain as a minor any of:

- ① infinitely many disjoint copies of either  $K_{3,3}$  or  $K_5$ ;
- ② infinitely many copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for a common vertex;
- ③ infinitely many copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for two common vertices; or
- ④  $K_{3,\aleph_0}$ .

# Our result: embeddability in bounded genus

Theorem (Christian, Richter, and S., 2011<sup>+</sup>)

A countable graph  $G$  embeds in some (orientable) surface of bounded genus if and only if  $G$  does not contain as a minor any of:

- ① infinitely many disjoint copies of either  $K_{3,3}$  or  $K_5$ ;
- ② infinitely many copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for a common vertex;
- ③ infinitely many copies of either  $K_{3,3}$  or  $K_5$  that are disjoint except for two common vertices; or
- ④  $K_{3,\aleph_0}$ .

A (slightly surprising?) consequence

There is no distinction between embeddability in some orientable surface and embeddability in some surface. In other words, no graph can embed in some (non-orientable) surface and have arbitrarily large projective grids.

# The interesting direction

The “only if” part is easy: a graph with infinitely many (sufficiently disjoint) copies of  $K_{3,3}$  or  $K_5$ , or with  $K_{3,\aleph_0}$ , cannot be embedded in any surface of bounded genus.

The “if” part is the interesting one.

# Getting started

For the rest of the talk, for brevity,

“surface” means *bounded genus, orientable* surface.

# Getting started

For the rest of the talk, for brevity,

“surface” means bounded genus, orientable surface.

A graph is **good** if it can be embedded in some surface.

Otherwise it is **bad**.

A bad  $G$  has infinitely many disjoint copies of  $K_{3,3}$  or  $K_5$  or:

There is a  $J \subseteq G$ , and a vertex  $u_1$  of  $J$ , such that  $J$  is bad and  $J - u_1$  is good.

- 1 Let  $G_0 := G$ , and as long as  $G_i$  has a subgraph  $H_{i+1}$  (may choose finite, if one exists) that contracts to  $K_{3,3}$  or  $K_5$ , set  $G_{i+1} := G_i - V(H_{i+1})$ .

A bad  $G$  has infinitely many disjoint copies of  $K_{3,3}$  or  $K_5$  or:

There is a  $J \subseteq G$ , and a vertex  $u_1$  of  $J$ , such that  $J$  is bad and  $J - u_1$  is good.

- 1 Let  $G_0 := G$ , and as long as  $G_i$  has a subgraph  $H_{i+1}$  (may choose finite, if one exists) that contracts to  $K_{3,3}$  or  $K_5$ , set  $G_{i+1} := G_i - V(H_{i+1})$ .
- 2 If for every positive  $i$ ,  $H_i$  exists, then we are done ( $G$  contains as a minor infinitely many disjoint copies of either  $K_{3,3}$  or  $K_5$ ). Thus we may assume that for some  $i$ ,  $G_i$  has no Kuratowski minor; so  $G_i$  is planar. Note that  $G_i$  is obtained from  $G$  by the deletion of finitely many vertices  $v_1, v_2, \dots, v_k$ .


A bad  $G$  has infinitely many disjoint copies of  $K_{3,3}$  or  $K_5$  or:

There is a  $J \subseteq G$ , and a vertex  $u_1$  of  $J$ , such that  $J$  is bad and  $J - u_1$  is good.

- 1 Let  $G_0 := G$ , and as long as  $G_i$  has a subgraph  $H_{i+1}$  (may choose finite, if one exists) that contracts to  $K_{3,3}$  or  $K_5$ , set  $G_{i+1} := G_i - V(H_{i+1})$ .
- 2 If for every positive  $i$ ,  $H_i$  exists, then we are done ( $G$  contains as a minor infinitely many disjoint copies of either  $K_{3,3}$  or  $K_5$ ). Thus we may assume that for some  $i$ ,  $G_i$  has no Kuratowski minor; so  $G_i$  is planar. Note that  $G_i$  is obtained from  $G$  by the deletion of finitely many vertices  $v_1, v_2, \dots, v_k$ .
- 3 For  $j = 1, 2, \dots, k$ , consider  $G^j := G - \{v_{j+1}, v_{j+2}, \dots, v_k\}$ . There is a least  $j$  so that  $G^j$  does not embed in any surface. Set  $J_0 := G^j$ , and  $u_1 := v_j$ . Thus  $J_0$  is a subgraph of  $G$  that does not embed in any surface ( $J_0$  is bad), yet  $J_0 - u_1$  does ( $J_0 - u_1$  is good).




**REPEAT AND GET:** Either  $G$  has one of the listed minors or  $\exists$ :



$M \subseteq G$

- $M$  is bad
- $M - u_i$  is good for each  $i$
- $M - A$  has no subdivision of  $K_{1,3}$  with  $u_1, u_2, u_3$  as the degree 1 vertices.

**REPEAT AND GET:** Either  $G$  has one of the listed minors or  $\exists$ :




$$M \subseteq G$$

- $M$  is bad
- $M - u_i$  is good for each  $i$
- $M - A$  has no subdivision of  $K_{1,3}$  with  $u_1, u_2, u_3$  as the degree 1 vertices.

**IF THIS HAPPENS:** Every component of  $M - (A \cup \{u_1, u_2, u_3\})$  attaches to at most two of  $u_1, u_2, u_3$ . Let  $N_{j,k}$  be the subgraph of  $M$  induced by the vertices in  $A \cup \{u_j, u_k\}$  and all components of  $M - (A \cup \{u_1, u_2, u_3\})$  that attach to at most  $u_j$  and  $u_k$ .

**REPEAT AND GET:** Either  $G$  has one of the listed minors or  $\exists$ :



$M \subseteq G$

- $M$  is bad
- $M - u_i$  is good for each  $i$
- $M - A$  has no subdivision of  $K_{1,3}$  with  $u_1, u_2, u_3$  as the degree 1 vertices.

**IF THIS HAPPENS:** Every component of  $M - (A \cup \{u_1, u_2, u_3\})$  attaches to at most two of  $u_1, u_2, u_3$ . Let  $N_{j,k}$  be the subgraph of  $M$  induced by the vertices in  $A \cup \{u_j, u_k\}$  and all components of  $M - (A \cup \{u_1, u_2, u_3\})$  that attach to at most  $u_j$  and  $u_k$ .

Each of  $N_{1,2}$ ,  $N_{1,3}$ , and  $N_{2,3}$  embeds in some surface. . . combine these embeddings to obtain an embedding of  $M$  (contradiction!).

Continuing in the countably infinite theme...

*A countably infinite number of men went into a pub. The first one ordered a pint. The second ordered a half-pint. The third ordered a quarter of a pint ... The barkeeper, with a face full of disgust, finally poured two pints and put them on the bar and said, "It's good when people know their limits."*

Continuing in the countably infinite theme...

*A countably infinite number of men went into a pub. The first one ordered a pint. The second ordered a half-pint. The third ordered a quarter of a pint ... The barkeeper, with a face full of disgust, finally poured two pints and put them on the bar and said, "It's good when people know their limits."*

*Thanks for your attention!*